

# DuPont fungicide program recommendations

*More complementary chemistries ... for broad-spectrum protection you can count on*



Recommended fungicide programs — Spray every 5 to 7 days			
CROP	TOMATOES	PEPPERS	CUCURBITS
Application 1	Spray DuPont™ Tanos™ (8 oz/acre) plus Manzate® + Kocide® 2000 (ManKocide® can be substituted for Manzate® + Kocide® tank mixture)	Spray Tanos™ (8 oz/acre) plus Kocide® 2000 + Manex® <b>or</b> Tanos™ 10 oz/acre plus Kocide® 2000	Spray Tanos™ (8 oz/acre) plus Manzate® (plus Kocide® 2000 if bacterial fruit blotch is a concern)
Application 2	Spray Manzate® plus Kocide® 2000 (or Kocide® 2000 plus chlorothalonil if target spot is present)	Spray Manex® plus Kocide® 2000	Spray Manzate® / ManKocide® (plus DuPont™ Curzate® 60DF if downy mildew is a concern) <b>or</b> chlorothalonil (if powdery mildew is a concern)
Application 3	Spray Tanos™ (8 oz/acre) plus Manzate® + Kocide® 2000 (ManKocide® can be substituted for Manzate® + Kocide® tank mixture)	Spray Tanos™ 8 oz/acre plus Kocide® 2000 + Manex® <b>or</b> Tanos™ 10 oz/acre plus Kocide® 2000	Spray a strobilurin plus Kocide®
Application 4	Spray Manzate® plus Kocide® 2000 (or Kocide® 2000 plus chlorothalonil if target spot is present)	Spray Manex® plus Kocide® 2000	Spray Manzate® / ManKocide® (plus Curzate® 60DF if downy mildew is a concern) <b>or</b> chlorothalonil (if powdery mildew is a concern)
Application 5	Repeat application 1	Repeat application 1	Repeat application 1
Comments	* Use Kocide® 2000 only if bacterial diseases are present. * This program has 4 modes of action to manage fungi and bacteria.	* This program has 4 modes of action to manage fungi and bacteria.	* Include other fungicides in program for gummy stem blight, belly rot and powdery mildew control. * This program has 5 modes of action to manage fungi and bacteria. * Do not make more than four (4) applications of Tanos™ or other Group 11 fungicides (all strobilurins or fenamidone) per crop per acre per season.

For more information visit our Web site at [tanos.dupont.com](http://tanos.dupont.com)

## Precautions

- These tank-mix programs must be sprayed in numerical order throughout the crop.
- Do not make more than one application of Tanos™ before alternating with a fungicide that has a different mode of action.<sup>1</sup>

<sup>1</sup> Tanos™ must be applied in tank mix and in alternation with appropriate contact fungicides which have a different mode of action: EBDCs, such as Manzate®, copper, such as Kocide® 2000 and chlorothalonil are recommended. Do not tank mix or alternate with fungicides from EPA's Target Site of Action Group 11 (includes all strobilurins, Famoxate™ and fenamidone).

*This reference guide is not intended as a substitute for the product label for the products referenced herein. Product labels for the above products contain important precautions, directions for use and product warranty and liability limitations that must be read before using the product. Applicators must be in possession of the product label(s) at the time of application. Always read and follow all label directions and precautions for use when using any pesticide alone or in tank mix combinations.*

The DuPont Oval, DuPont™, The miracles of science™, Curzate®, Famoxate™, Kocide®, Manex®, ManKocide®, Manzate® and Tanos™ are registered trademarks or trademarks of DuPont or its affiliates.

Copyright © 2001-2004 E.I. du Pont de Nemours and Company. All Rights Reserved. 10/04  
Reorder No.: K-05878 (Replaces K-02361)



The miracles of science™