

# DuPont™ Lannate® LV

insecticide

*Lannate® LV insecticide quickly controls armyworm, fall armyworm, beet armyworm, sorghum midge, corn earworm and sorghum webworm*

DuPont™ Lannate® LV is an effective tool for pest control in grain sorghum. The key to controlling sorghum pests is to regularly scout your fields and to properly time your Lannate® LV insecticide applications when it is economical to spray. When scouting your sorghum\*, here's what to look for:

- Fall armyworms deep in the whorl before evidence of damage to leaves.
- Evidence of feeding on leaves by armyworms.
- Sorghum midges early in the morning or late in the day when they are most active; or, bag the blooming heads and shake, to determine an infestation of adult flies.
- Evidence of worms feeding on heads in flower until hard dough stage. Look for webworms, especially in late-planted fields.
- When you decide it's time to spray Lannate® LV, follow this program:

## Early season

### Fall armyworm, corn earworm

- For these budworms, apply  $\frac{3}{4}$  to 1.5 pints of Lannate® LV per acre when larvae are found in the whorl. Make additional treatments as needed.

### Armyworm, beet armyworm

- For these leaf feeders, apply  $\frac{3}{4}$  to 1.5 pints of Lannate® LV per acre when significant defoliation of plants is detected. Repeat as needed.

## Mid-to-late season

### Sorghum midge

- Use  $\frac{3}{4}$  to 1.5 pints of Lannate® LV per acre. Apply when 50 percent of the plants have begun to bloom. If adults are active three to five days after treatment, make additional treatments as needed, for midges frequently reinfest treated fields.

### Sorghum webworm

- Apply 1.5 pints of Lannate® LV per acre when larvae reach economic thresholds. Repeat as needed.



### Corn earworm, fall armyworm

- Apply  $\frac{3}{4}$  to 1.5 pints of Lannate® LV per acre when worms appear on head in damaging numbers. Repeat as needed.

### Armyworm, beet armyworm

- Apply  $\frac{3}{4}$  to 1.5 pints of Lannate® LV per acre when significant defoliation of plants is detected.

Important: Use sufficient spray volume (see label for ground and air recommendations) to obtain uniform coverage. Do not apply within 14 days of harvest. Do not graze or feed treated plants to livestock within 14 days of application.

## For more information

Please contact your local DuPont crop protection retailer or DuPont representative to learn more about controlling sorghum pests with Lannate® LV insecticide. And visit us on the Web at [lannatelv.dupont.com](http://lannatelv.dupont.com)

\* Not for use on sweet sorghum.

Lannate® LV is a restricted-use pesticide.

*This reference guide is not intended as a substitute for the product label for the product(s) referenced herein. Product labels for the above product(s) contain important precautions, directions for use and product warranty and liability limitations that must be read before using the product. Applicators must be in possession of the product label(s) at the time of application. Always read and follow all label directions and precautions for use when using any pesticide alone or in tank mix combinations.*

*The DuPont Oval Logo, DuPont™, The miracles of science™ and Lannate® are trademarks or registered trademarks of DuPont or its affiliates.*

*Copyright © 2005-2009 E.I. du Pont de Nemours and Company. All Rights Reserved. 1/09 Reorder No.: K-15030 (Replaces K-06213)*



*The miracles of science™*